

Lakeshore West Dental

3390 Lake Shore Blvd., W.
Etobicoke, Ontario, M8W 1M9

Office: 416-251-5707
lakeshorewestdental@gmail.com

SELF CARE AFTER IMPLANT SURGERY

BLEEDING

- Some blood may ooze from the surgery area for a day or two, which is normal. If bleeding continues, contact the office.

WOUND CARE

- **DO NOT SMOKE** for at least 7 days because it promotes infection, bleeding and interferes with healing.
- **DO NOT SPIT** or suck through a straw, since this promotes bleeding.

DISCOMFORT

- Some discomfort is normal after surgery but can be controlled by medication prescribed by the dentist.
- Follow the prescription as advised by your dentist and the directions written on the container. If the medication is not strong enough please contact the office.

DIET

- It is important to drink plenty of fluids (avoid alcohol) and eat regular meals the day after surgery. Soft food such as pudding, yogurt, ice cream, mashed potatoes and scrambled eggs may be the most comfortable to start with. No solids the day of surgery.

ORAL HYGIENE

- **DO NOT RINSE WITH ANY TYPE OF LIQUID** vigorously the day of surgery. If sutures have been placed rinsing may cause premature loss of sutures.
- Continue to brush your teeth but avoid the area of surgery and do not use toothpaste after the surgery. After brushing simply allow the water to fall from your mouth, do not spit vigorously.

SWELLING

- Swelling after surgery is a normal body reaction. It reaches its maximum 3 days after surgery and usually lasts 4 to 6 days. Intraoral ice application will reduce swelling and pain and is advised for at least 3 days post operatively.

BRUISING

- You may experience some mild bruising around the surgical area. This response is normal for some people and should not cause you any alarm. The bruising will disappear in a week or two.

STIFFNESS

- After surgery you may experience jaw muscle stiffness and limited opening of your mouth, this is normal and should improve in 5 to 10 days.

STITCHES

- If stitches have been placed, you MAY need to have them removed in 7 to 14 days and an appointment should be made.
- Dissolving sutures may have been used and usually take a week to disappear.

REST

- Avoid strenuous activity on the day of surgery.

PROBLEMS

- If you have any problems or questions please call the office as soon as possible or the Dental Emergency Clinic in your region.
- ***Or Dr. Sam Patel at (416) 251-5707 or (416) 570-5707.***